

AD-A230 892

DTIC FILE COPY

①

REPORT DOCUMENTATION PAGE			
1. Agency Use Only.		2. Report Date. June 4-7, 1989	3. Type of Report and Dates Covered. Conference June 4-7, 1989
4. Title and Subtitle. Vector EOF Analysis of SSH and Wind Stress for the Geosat Pre-Erm Mission in the Gulf of Mexico			5. Funding Numbers. Program Element No. 63704N Project No. 0101 Task No. 0101 Accession No. DN258117
6. Author(s). Don Johnson			
7. Performing Organization Name(s) and Address(es). Naval Ocean Research and Development Activity Code 321 Ocean Science Directorate, Stennis Space Center, MS 39529			8. Performing Organization Report No. AB 89:321:044
9. Sponsoring/Monitoring Agency Name(s) and Address(es). SPAWAR			10. Sponsoring/Monitoring Agency Report Number.
11. Supplementary Notes.			
12a. Distribution Availability Statement. Approved for public release; Distribution is unlimited.			12b. Distribution Code.
13. Abstract (Maximum 200 Words). See Text			
14. Subject Terms. EOF, Gulf of Mexico, GEOSAT, SSH, Wind Stress			15. Number of Pages. 1
17a. Security Classification. (of Report) Unclassified			16. Price Code.
17b. Security Classification. (of Page)			17c. Security Classification. (of Abstract) Unclassified
			18. Media Type.

SF 298 (2/89)

DTIC
ELECTE
JAN 31 1991
S E D

91 1 30 023

Vector EOF Analysis of SSH and Wind Stress for
the GEOSAT Pre-ERM mission in the Gulf of
Mexico.

D. R. Johnson (Naval Ocean Research and
Development Activity, Code 321, SSC, MS
39529; 601-688-4691; Omnet TMail D.Johnson

Sea Surface Height (SSH) data collected during
the Pre-exact Repeat Mission of the U.S. Navy's
GEOSAT satellite were analyzed together with
wind stress data from FNOC to determine basin
scale patterns of coherent variability. Time
series of SSH variations were formed from
altimeter cross-over points in 3-degree
diamond-shaped grids in the Gulf of Mexico
(GOM). Similarly, wind stress time series were
formed at selected grid points in the GOM and
in the North Equatorial Trade wind region of
the Atlantic.

The two sets of time series, one year in
length, were subjected to Empirical Orthogonal
Function analysis. The first EOF mode
contained more than twice the variance of the
second mode, with the largest amplitudes
(eigenvectors) of the pattern occurring just
west of the loop current intrusion for SSH and
in the western and northern Gulf for wind
stress. A comparison is made with similar
combined data for the Northern Indian Ocean
where ocean response to Monsoon winds is
strong, and better understood. In addition to
the EOF analysis, events are followed through
the Gulf in both wind and SSH records.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	